

# 'Study of the nutritional characteristics of previously analysed edible mushrooms as biofertilisers'

Enhancing nutritious food availability through promotion of native edible tree/shrub species in Sub-Saharan Africa



The present project will be carried out in Benin, Burkina Faso, Mali and Niger (Sub-Saharan Africa). This region is characterized by high population growth, unpredictable food provision and high poverty rates. This situation is already and will further be worsened by climate change effects which affect agricultural production.

## OBJECTIFS

'From a collection of fungal strains isolated in Ghana and representing the genera *Lentinus*, *Pleurotus* and *Ganoderma*, the objectives of this study were the following :  
To highlight certain physiological properties likely to be involved in the biological mechanisms promoting plant growth  
To demonstrate under controlled conditions the effect of inoculation of the culture substrate by these strains on the growth of 2 plant species, tomato and durum wheat'



## ACTIONS

'The fungal strains (7) were maintained on a nutrient medium of the "L-Broth" type and were subcultured on selective media in order to demonstrate the enzymatic activities of the esterase, chitinase and phosphatase types as well as their capacity to produce siderophores.  
In a second phase, these strains were multiplied on a vermiculite substrate moistened with a nutrient medium of the "L-Broth" type. When the substrate was completely colonised by the fungal isolates, the inoculum produced was mixed with a culture substrate (atapulgit) where young tomato or durum wheat seedlings were planted.'

## RESULTATS

'All strains tested have the ability to produce esterase enzymes, whereas only isolates from the genus *Lentinus* show chitinolytic activity. The production of siderophores was detected in strains belonging to the genera *Ganoderma* and *Lentinus* and in one strain of *Pleurotus* (strain PTR-K). The promoting effects of strain inoculation were also measured to varying degrees depending on the isolate in tomato after 6 weeks in the greenhouse  
Fungal inoculation stimulated the growth of young durum wheat seedlings after one month of cultivation in the greenhouse with a maximum effect obtained with the SqW strain (*Lentinus* sp.) (Figure below).'

## PERSPECTIVES

'The perspectives of this work will mainly consist in optimising the production of mushrooms AND their biofertilising properties by  
Testing a greater diversity of edible fungi  
Manipulating the composition of the culture substrate  
The aim of the project will be to identify high-performance strains which, depending on an appropriate multiplication technique, will be able to offer significant production capacity but also proven biofertilising properties.'

**Responsable :** Dr. Amadou M. Kouyaté  
kouyate01@hotmail.com

**Etablissement :** AgroParisTech, CIHEAM IAMM

**Unite de recherche :** AGAP, AIDA

**Pays concernés:** Afrique du Sud

**Date de démarrage :** 07/02/2016

**Date de clôture :** 07/05/2016

**Montant :** 500 000 €

